

Descriptions of Map Units

COASTAL PLAIN

- Fill: Light-colored deposits of natural road material, including gravel, used to extend those land and/or to fill a low-lying area such as where a road crosses a valley or marsh.
Shoreline Deposits (Holocene): Beach and dune deposits along the shoreline of Delaware Bay.
Alluvial Deposits (Holocene): Browns, light yellow-orange, and gray fine to coarse quartz sand, silt, clay, and fine to medium gravel.
Swamp Deposits (Holocene): Structureless, black to brown, organic-rich, silty and clayey, fine to coarse quartz sand with thin interbeds of medium to coarse quartz sand.
Marsh Deposits (Holocene): Structureless to finely laminated, black to dark-gray, organic-rich silty clay to clayey silt with discontinuous beds of peat and rare shells.
Alluvium and Swamp Deposits (upper Pleistocene to Holocene): Found along the headwaters of streams that drain into Chesapeake Bay.
Undrained Depression Deposits (upper Pleistocene to lower Holocene): A belt of upland depressions stretches across southern New Castle and northern Kent counties.
Seaside Cores Formation (upper Pleistocene): Heterogeneous unit of light gray to brown to light yellowish-brown, coarse to fine sand, gravelly sand and pebbly gravel.
Lynch Heights Formation (upper Pleistocene): Heterogeneous unit of light gray to brown to light yellowish-brown, medium to fine sand with discontinuous beds of coarse sand, gravel, silt, fine to very fine sand, and organic-rich clayey silt to silty sand.
Turtle Branch Formation (informal unit) (upper Pleistocene): One to five feet of gray coarse sand and pebbles overlain by one to ten feet of tan to gray clayey silt to silty clay.
Columbia Formation (middle Pleistocene): Yellowish to reddish-brown, fine to coarse, idiosyncratic quartz sand with varying amounts of gravel.
Beaverdam Formation (upper Pliocene): Light gray to white coarse to very coarse sand with beds of fine to medium sand.

CROSS-SECTION UNITS (not shown on map)

- St. Marys Formation (upper Miocene) subsurface only: Light gray to blue gray, fine to medium, silty sand and clayey silt.
Choptank Formation (middle to upper Miocene) primarily subsurface: Gray to grayish-brown, clayey silt to silty clay interbedded with gray to light gray silt to fine to coarse quartz sand.
Calvert Formation (lower to middle Miocene) primarily subsurface: Gray to grayish-brown, clayey silt to silty clay interbedded with gray to light gray silt to fine to coarse quartz sand.
Piney Point Formation (upper Eocene) subsurface only: Bright green, fine to coarse, shelly, glauconitic (20 to 40% glauconite), quartz sand. Silty and clayey toward the bottom and coarser upward.

DISCUSSION OF MAP


This map shows the surficial geology of Kent County, Delaware at a scale of 1:100,000. Maps at this scale are useful for viewing the general geologic framework on a county-wide basis, determining the geology of watersheds, and recognizing the relationship of geology to regional or county-wide environmental or land-use issues. This map, when combined with the subsurface geologic information, provides a basis for locating water supplies, mapping ground-water recharge areas, and protecting ground and surface water.

Kent County is located within the Atlantic Coastal Plain which is composed of seaward-dipping strata of sand, silt, and clay. The surficial geology consists of units ranging in age from late Tertiary (Pliocene) to Holocene (swamp, marsh, and alluvium). The surficial units are primarily composed of sand. Differentiation between surficial units and separation from underlying, older sand bodies is not always possible.




The oldest surficial unit of any great extent in Kent County is the Beaverdam Formation. The Beaverdam represents a major period of fluvial deposition in the region that followed a period of erosion in the late Miocene. The Beaverdam rests on an unconformity that truncates progressively older units one goes from south to north in Kent County (cross section B-B', C-C').

After deposition of the Beaverdam, the first major glacialiations occurred in northeastern North America during the early Pliocene. As these glacial melt, huge volumes of meltwater flowed from the headwaters of the Delaware and Susquehanna rivers eroding away much of the Beaverdam Formation which is now southern New Castle County. These meltwater rivers transported large volumes of sediment that were deposited as the Columbia Formation.

Since the deposition of the Columbia, sea level has risen and fallen several times. During interglacial high stands sea level, at about 225,000 and 100,000 years ago, the Lynch Heights and Seaside Cores Formations, respectively, were deposited on the margins of an ancestral Delaware Bay (Ramsey, 1997).



Correlation of Geologic Units



DISCUSSION (cont.)

During the glacial periods of the Pleistocene, Delaware was subjected to cold climate and periglacial conditions that modified the landscape with cold-climate features such as unindented depressions, Carolina Bays, and alluvium and swamp deposits. Andes and Howard (1999) documented the association of freshwater features in the subsurface with the surficial unindented depressions found in southern New Castle County and northeastern Kent County.

The rise of sea level during the past 12,000 years has resulted in the development of the Holocene swamps, marshes, and shoreline deposits. On the uplands, ponding in undrained depressions, upland swamps, and Carolina Bays contributes to the present disposition of fine-grained, organic swamp sediments.

DISCUSSION OF CROSS SECTIONS

Three cross sections show the relationship of surficial and subsurface stratigraphic units. Section A-A' and B-B' which follow closely the same path as sections E-E' and S-S' of McLaughlin and Velz (2006), show the principal aquifers of Kent County. Section C-C' is a west-east section showing the stratigraphic units in the southern portion of Kent County, which extends section A-A' of Ramsey (1997) to the west.

Surface topography of the cross sections was constructed by taking a data slice along the line of cross-section from the digital elevation model (DEM) using Surfer software. Correlation of stratigraphic units was done by using geophysical logs, lithologic descriptions, and geologists' logs, and examination of samples. Correlation within the cross-sections is a depth which shows the base of the Calvert Formation. The Calvert Formation in northern and central Kent County lies beneath the surficial deposits.

Recognition of Quaternary units presents a challenge because all of the units are predominantly sand, and units also commonly overlie sands of the Miocene Choptank and Calvert Formations that are normally similar to the Quaternary sands. Use of gamma log response offers only a limited tool for separation of the sands. Subdrilling Miocene sands were correlated from down dip where the sands are bounded by fine grained clay or shale.

Surface topography shown on the cross sections also shows the relationship between the land surface and underlying surficial units. Cross-section A-A' shows the cut and fill relationship between the Columbia Formation and younger units including the Lynch Heights and Seaside Cores Formations.

REFERENCES

Andes, A. S. and Howard, C. S., 1998. Analysis of deformation features at the Pollack Farm Site, Delaware. In: Benson, R.N., ed. Geology and paleontology of the lower Miocene Pollack Farm Fossil Site, Delaware: Delaware Geological Survey Special Publication No. 21, p. 47-53.
Andes, A.S. and Ramsey, K.W., 1995a. Geologic Survey of the Seaford area, Delaware: Delaware Geological Survey Geologic Map Series No. 9, scale 1:24,000.
Andes, A.S. and Ramsey, K.W., 1995b. Geologic Survey of the Seaford area, Delaware: Delaware Geological Survey Geologic Map Series No. 5, scale 1:24,000.
Benson, R.N., 1996. Stratigraphy of the post-Potomac Cretaceous-Tertiary rocks of Central Delaware: Delaware Geological Survey Bulletin No. 20, 28 p.
Benson, R.N., Jordan, R.E., and Spangler, N., 1985. Geological study of the Cretaceous and Tertiary Section, Test Well J632-04, Central Delaware: Delaware Geological Survey Bulletin No. 17, 69 p.
Benson, R.N. and Pickett, T.E., 1986. Geology of south-central Kent County, Delaware: Delaware Geological Survey Geologic Map Series No. 7, scale 1:24,000.
Groot, J.J. and Jordan, R.E., 1999. The Pliocene and Quaternary geology of Delaware: Delaware Geological Survey Report of Investigations No. 58, 30 p.
Jordan, R.E., 1964. Columbia (Pliocene) deposits of Delaware: Delaware Geological Survey Bulletin No. 12, 69 p.
Mathews, E.D. and Ireland, W., Jr., 1971. Soil Survey of Kent County, Delaware: U.S. Department of Agriculture, 66 p.
McLaughlin, P.P. and Velz, C.C., 2006. Geology and extent of the confined aquifers of Kent County, Delaware: Delaware Geological Survey Report of Investigations No. 72, 40 p.
Pickett, T.E. and Benson, R.N., 1977. Geology of the Smyrna-Clayton area, Delaware: Delaware Geological Survey Geologic Map Series No. 5, scale 1:24,000.
Ramsey, K.W., 1996. Geology of the Dover area, Delaware: Delaware Geological Survey Geologic Map Series No. 6, scale 1:24,000.
Ramsey, K.W., 1992. Coastal response to late Pliocene climate change, Middle Atlantic Coastal Plain, Virginia and Delaware. In: Pickett, J.H. III and Wolinsky, J.E., eds. Quaternary Coasts of the United States: marine and lacustrine systems, SEPM Special Publication No. 48, p. 121-127.
Ramsey, K.W., 1992. Coastal response to late Pliocene climate change, Middle Atlantic Coastal Plain, Virginia and Delaware. In: Pickett, J.H. III and Wolinsky, J.E., eds. Quaternary Coasts of the United States: marine and lacustrine systems, SEPM Special Publication No. 48, p. 121-127.
Ramsey, K.W., 1996. Radiocarbon dates from Delaware: a compilation. Delaware Geological Survey Report of Investigations No. 54, 18 p.
Ramsey, K.W., 1997. Geology of the Milford and Millsboro Quadrangles, Delaware: Delaware Geological Survey Report of Investigations No. 55, 40 p.
2001. Geologic Map of the Hellenburg and Millsboro Quadrangles, Delaware: Delaware Geological Survey Geologic Map Series No. 11, scale 1:24,000.
2003. Geologic Map of the Croves and Cape Henlopen Quadrangles, Delaware: Delaware Geological Survey Geologic Map Series No. 12, scale 1:24,000.
2006. Geology of New Castle County, Delaware: Delaware Geological Survey Geologic Map Series No. 13, scale 1:100,000.
Webb, R.S., 1990. Late Quaternary water-level fluctuations in the northeastern United States: Ph. D. Dissertation, Brown University, Providence, R.I., 131 p.

Geologic Map of Kent County, Delaware

by
Kelvin W. Ramsey

2007

MAP CREDITS
Base Map: Universal Transverse Mercator, Zone 18
North American Datum of 1983 (NAD83)
USGS Delaware Hydrography Lines 2002.
USGS Delaware State Boundary Lines 2002.
USGS Delaware Boundaries - County Boundary Lines 2002.
The Delaware Office of State Planning, Coordination, Delaware Management, 2005
Delaware Department of Transportation Centerline for Delaware, 2006
Kent DEM: Dataserver resolution, http://www.sde.state.de.us/DEM/arcserve/arcserve.html
Other: Cartography by William T. Wang, Delaware Geological Survey
Edited by Stefanie J. Basso, Delaware Geological Survey
Map layout and design by Allison T. Wang and Stefanie J. Basso
Acknowledgements: This work was funded in part by the cooperative agreement between the Association of American State Geologists and the U.S. Geological Survey under STATEMAP Program Grants 04HQPA003.